

ENERMAX®

ENERGETIC BIOSTIMULANT TO INCREASE AND STANDARDIZE FRUIT SIZE

- Directs photo-assimilates and nutrients selectively to the fruit
- Promotes cell division- and stimulates the synthesis of new proteins.
- Prevents stress situations and or deformation of the fruit.

ENERMAX Confirms its potential for:

- Earlier fruit ripening.
- 20% pack out in the first picking compared to control.
- Increases size of the weakest & smallest fruit or tubers.
- Results in uniform even distribution of fruit & tuber size.
- Improves fruit quality and brix.

APPLICATION TIMING:

CROP	First Application	Second Application
Avocado	Fruit size:25-30mm diameter	40-30 days prior to harvest.
Citrus		
Blueberries	Fruit size: 2-5mm diameter	14 days later
Macadamia	Nut size: 5-10mm diameter	14 days later
Potato	Tuber initiation	Tuber bulking.
Tomato	Second truss set	14-21 days later

APPLICATION RATES & RECOMMENDATIONS:

Crop	Foliar (g/100L)	Soil (kg/ha)	Recommendations
Apricots Peaches Plums	50-100 g	0.5-1.5 kg	Apply in 500-1500l/ha of water when fruits are 10-15mm in diameter. Repeat at 40-30 days prior to harvest.
Avocado	50-100 g	0.5-1.5 kg	Apply in 500-1500l/ha of water when fruits are 20-30mm in diameter. Repeat at 40-30 days prior to harvest.
Banana	50-100 g	0.5-1.5 kg	Apply in 300-600l/ha of water when bunches begin to emerge and repeat every 14-21 days.
Blueberries	50-100 g	0.5-1.5 kg	Apply in 500-1500l/ha of water when fruits are 2-5mm in diameter. Repeat at 14-21 days later.
Cabbage	50-100 g	0.5-1.5 kg	Apply in 300-600l/ha of water when head begin to emerge and repeat 14-21 days later.
Carrot	50-100 g	0.5-1.5 kg	Apply in 300-600l/ha of water when tubers begin to bulk and repeat 14-21 days later
Cauliflower Broccoli	50-100 g	0.5-1.5 kg	Apply in 300-600l/ha of water when head is 50 mm in diameter and repeat 14 days later.
Chillies Paprika Peppers	50-100 g	0.5-1.5 kg	Apply in 300-600l/ha of water at fruit set and repeat every 14-21 days to a maximum of 5 sprays per growing season.
Citrus	50-100 g	0.5-1.5 kg	Apply in 500-1500l/ha of water when fruits are 20-30mm in diameter. Repeat 14 days later and 40-30 days prior to harvest.
Coffee	50-100 g	0.5-1.5 kg	Apply in 500-1500l/ha of water at first berry set. Repeat at 14-21 days later and 40-30 days prior to harvest.
Cucumbers Pumpkins Melons Squash	50-100 g	0.5-1.5 kg	Apply in 300-600l/ha of water at fruit set and repeat every 14-21 days to a maximum of 5 sprays per growing season.
Grapes	50-100 g	0.5-1.5 kg	Apply in 500-1500l/ha of water when fruits are 5mm in length. Repeat at 14 days and 28 days after the first spray.
Granadilla	50-100 g	0.5-1.5 kg	Apply in 500-1500l/ha of water at fruit set. Repeat every 14-21 days.
Groundnuts			Apply in 300-600l/ha of water at pod set. Repeat at 14 days and 30 days after the first spray.
Litchi	50-100 g	0.5-1.5 kg	Apply in 500-1500l/ha of water when fruit is 5-10 mm length. Repeat at 14 days and 28 days after the first spray.
Macadamia	50-100 g	0.5-1.5 kg	Apply in 500-1500l/ha of water when fruits are 10-15mm in diameter. Repeat at 14 days and 40-30 days prior to harvest.
Mango	50-100 g	0.5-1.5 kg	Apply in 500-1500l/ha of water when fruits are 5-10mm in diameter. Repeat at 14 days and 40-30 days prior to harvest.
Maize	50-100 g	0.5-1.5 kg	Apply in 300-600l/ha of water at 5-7 leaf stage. Repeat between silking and grain setting stage.
Onions	50-100 g	0.5-1.5 kg	Apply in 500-1500l/ha of water when tubers are 10-15mm in diameter. Repeat 40-30 days prior to harvest.
Papaya	50-100 g	0.5-1.5 kg	Apply in 500-1500l/ha of water at fruit set. Repeat every 14-28 days.
Peas	50-100 g	0.5-1.5 kg	Apply in 300-600l/ha of water at pod set. Repeat at 14 days later.
Pecan	50-100 g	0.5-1.5 kg	Apply in 500-1500l/ha of water when fruits are 10-15mm in diameter. Repeat at 14 days and 28 days after first spray.
Pineapple	50-100 g	0.5-1.5 kg	Apply in 500-1500l/ha of water 30 days after emergence of flower head. Repeat every at 14-28 days until 30 days prior to harvest.
Potatoes	50-100 g	0.5-1.5 kg	Apply in 300-600l/ha of water at tuber initiation. Repeat at tuber set and tuber bulking.
Soyabean	50-100 g	0.5-1.5 kg	Apply in 300-600l/ha of water at pod set. Repeat at 14 days later.
Sugar beans	50-100 g	0.5-1.5 kg	Apply in 300-600l/ha of water at pod set. Repeat at 14 days later.
Sweet Potato	50-100 g	0.5-1.5 kg	Apply in 300-600l/ha of water at tuber initiation. Repeat at tuber set and tuber bulking.
Tomatoes	50-100 g	0.5-1.5 kg	Apply in 300-600l/ha of water at grain set. Repeat every 14-21 days.
Wheat/Barley	50-100 g	0.5-1.5 kg	Apply in 300-600l/ha of water at grain set.

Contact: Robert Mazuru / Craig Chittenden / Dzingo Mafuvadze / Golden Zenda

Mobile: 0712 204 946 - 7 / 0773 212 051

Email: mazuru@pivotalagro.com / craig@pivotalagro.com / dzingo@pivotalagro.com