



Focus® Ultra



Chemical Group: cyclohexenone

A selective post emergent systemic herbicide for use in broadleaf crops against annual and perennial grasses and volunteer cereals.

FOCUS ULTRA can be used on the following crops –

Citrus, Cucurbits, Deciduous fruits, Dry beans, Forestry crops, Grape vines, Green beans, Groundnuts, Lucerne, Onions, Pineapples, Potatoes, Soybeans, Sunflowers, Tomatoes and Subtropical fruits including Avocados, Litchis, Macadamia nuts, Mangoes, Papayas and Pecan nuts.

	FOCUS ULTRA Rate in l / ha	
Weeds Controlled	Seedlings up to just before tillering	Tillering – heading
<i>Digitaria spp</i>	0.8	1.6
<i>Eleusine indica</i>	0.8	1.6
<i>Eragrostis spp.</i>	0.8	1.6
<i>Paspalum paspalodes</i>	0.8	2.0
<i>Panicum schinzii</i>	0.8	2.0
<i>Urochloa panicoides</i>	0.8	1.0
<i>Eragrostis spp.</i>	1.0	2.0
<i>Phalaris canariensis</i>	1.0	3.0
<i>Setaria spp.</i>	1.2	1.6
<i>Volunteer Wheat</i>	1.2	3.0
<i>Volunteer Maize</i>	1.2	2.0
<i>Shamva Grass (Rottboellia cochinchinensis)</i>	1.6	2.0
<i>Annual grasses (3 – 7 leaf stage)</i>	1.4	1.8
<i>Couch Grass (Cynodon dactylon)</i>	2.0	4.0

Application details:

- 1) FOCUS ULTRA must be applied post-emergence to target species, which should be actively growing at the time of application.
- 2) Allow at least 3 days between a FOCUS ULTRA application and any other herbicide treatment.
- 3) FOCUS ULTRA is formulated with an adjuvant and therefore additional spray adjuvants are not required.
- 4) The product is normally applied in a volume of 150-200 liters of water / ha
- 5) Weeds which have not emerged at the time of application will not be controlled.
- 6) FOCUS ULTRA can be safely applied over the top of broadleaf crops at any growth stage.
- 7) FOCUS ULTRA is rainfast one hour after application.
- 8) Grasses suffering from drought stress will not be adequately controlled.

Manufactured by **BASF**
Registered By: Pivotal Agro-Services (Private) Ltd.