



PIVOTAL
AGRO-SERVICES



PLANT HEALTH
CARE

Myconate®

A natural compound that stimulates the development of root Mycorrhizae Fungi and Phosphate Solubilizing Bacteria for improved root growth and tolerance to soil pathogens and environmental stress.

Trade Name: MYCONATE
Active Ingredients: Formononectin
Registration No: 18-G-29-1
Target crops & Pests: See Below Table

BENEFITS OF MYCONATE:

- ✓ **Early root establishment & growth**
- ✓ **Increased water availability & uptake**
- ✓ **Improved nutrient uptake**
- ✓ **Improved pathogen resistance**
- ✓ **Improved germination rates**
- ✓ **Uniform germination**
- ✓ **Tolerance to root diseases**
- ✓ **Better tolerance to drought stress**
- ✓ **Tolerance to soil salinity**

APPLICATION RECOMMENDATIONS

CROP	RATE/HA	APPLY AS SOIL DRENCH, SEED DRESSING OR SPRAY APPLICATION
Banana	50g	Apply to young plants after transplant and repeat every 3-4 months.
Citrus	50-75g	Apply first application at beginning of new root flush before start of season and second application at fruit set.
Row Crops	25-35g	Apply as seed dressing or in planting furrow before covering.
Macadamia	50-75g	Apply at the beginning of before the new season and second application at fruit set.
New Trees/ Vineyard Plantings	50g	Apply in a solution in the planting hole at planting and follow-up 4-6 weeks later.
Ornamentals Roses	50g	Apply as soil drench during root flush.
Pecan Nuts	50g	Apply 3-4 weeks after bud break when new signs of leaf growth appears.
Pome/Stone Fruit	50-75g	Apply 3-4 weeks after bud break when new signs of leaf growth appears.
Potatoes	50-75g	Apply directly to tubers before planting or in the planting furrow before covering the tubers.
Subtropical Fruit	50-75g	Apply 3-4 weeks after bud break when new signs of leaf growth appear.
Sugarcane	50-100g	Plant Cane: Apply directly onto laid plant sets before covering plant cane. Ratoon Cane: Apply 14-21 days after harvesting directly over the root zone.
Vegetables	50-75g	Apply once through irrigation at transplanting.
Vineyards	50g	Apply 3-4 weeks after bud break when new signs of leaf growth appear.
Wheat / Barley	35-50g	Apply as seed dressing at planting.

Contact: Robert Mazuru / Craig Chittenden / Dzingo Mafuvadze/ Golden Zenda

Mobile: 0712 204 946 -7 / 0773 212 051

Email: mazuru@pivotalagro.com / craig@pivotalagro.com / dzingo@pivotalagro.com

