



# Prestige 38 SC®

**A suspension concentrate fungicide for the preventive control of various fungal diseases.**

**Trade Name:** Prestige 38 SC  
**Active Ingredients:** Pyraclostrobin 128g/l + Boscalid 252g/l  
**Registration No:** 19-B-88-7-B-90-3  
**Target Crops & Pests:** See Below Table

## APPLICATION RATES AND RECOMMENDATIONS

| Crop                                                            | Disease/Problem                                             | Rate/Ha    | Application                                                                                                                                                                                                                                                                                                                                          |
|-----------------------------------------------------------------|-------------------------------------------------------------|------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Black Berries<br>Blue Berries<br>Goose Berries<br>Straw Berries | Anthrachnose<br>Botrytis<br>Leaf Rust<br>Septoria Leaf Spot | 300 ml     | Apply in 300-500L of water per hectare at post-harvest flush before the rains. Repeat at 10 to 14-day intervals up to a maximum of 3 sprays per season. Rotate with another fungicide with a different mode of action.                                                                                                                               |
| Chillies<br>Paprika<br>Pepper                                   | Anthrachnose<br>Cercospora<br>Powdery Mildew                | 500 ml     | Apply in 300-500L of water per hectare as a preventative spray at 10 to 14-day intervals up to a maximum of 3 sprays. Use shorter intervals under heavy disease pressure.                                                                                                                                                                            |
| Broccoli<br>Cabbage<br>Cauliflower                              | Botrytis<br>Downy Mildew                                    | 800 ml     | Apply as preventative spray in 300-500 of water per hectare at 10 to 14-day intervals. Use shorter intervals in prolonged wet weather up to a maximum of 3 sprays per season.                                                                                                                                                                        |
| Cucumber<br>Pumpkin<br>Squash                                   | Alternaria<br>Powdery Mildew<br>Stem Blight                 | 800 ml     | Apply in 300-500 of water as a full cover spray at 10 to 14-day intervals up to maximum of 3 sprays. Rotate with another fungicide with a different mode of action.                                                                                                                                                                                  |
| Onions                                                          | Purple Blotch                                               | 300 ml     | Apply in 300-500L of water per hectare at 14-day intervals up to a maximum of 3 sprays per season. Rotate with fungicides with different mode of action to manage resistance.                                                                                                                                                                        |
| Ornamentals<br>Carnations<br>Roses                              | Downy Mildew<br>Black Spot<br>Powdery Mildew                | 650-800 MI | Apply in a preventative program in 500L of water/ha. Repeat every 7 to 10 days depending on rain and disease pressure. The shorter interval of 7 days should be used where disease pressure and/or rainfall is high. Apply 3 sequential applications of <b>PRESTIGE 38 SC</b> and after that use another fungicide from another class e.g. CUPERDEM. |

|                           |                                                               |            |                                                                                                                                                                                                                                                       |
|---------------------------|---------------------------------------------------------------|------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Macadamia                 | Athracnose<br>Botrytis<br>Blossom Blight<br>Husk Spot         | 800 ml     | Apply as a full cover spray in 1000L of water at beginning of flowering as a preventative program. Repeat at 14-day intervals up to a maximum of 3 sprays per season rotating with a fungicide of different mode of action.                           |
| Peas<br>Sugar beans       | Alternaria<br>Anthracnose<br>Ascochyta<br>Botrytis            | 500-800 ml | Apply in 300-500L of water as a preventative spray before onset of diseases at 7 to10-day intervals up to a maximum of 3 sprays per season. Use in rotation with a fungicide of different mode of action.                                             |
| Potatoes                  | Early Blight<br>Late Blight                                   | 300 ml     | Apply in 300-500L of water per hectare as a preventative spray before the onset of the disease at 14-day intervals up to a maximum of 3 sprays per season. Use in rotation with a curative fungicide if Late Blight has set in.                       |
| Tomatoes                  | Early Blight<br>Late Blight                                   | 300 ml     | Apply in 300-500L of water as a preventative spray before onset of diseases at 7 to10-day intervals up to a maximum of 3 sprays per season. Use in rotation with a curative fungicide if Late Blight has set in.                                      |
| Pome Fruit<br>Stone Fruit | Apple Scab<br>Blossom Blight<br>Powdery Mildew<br>Sooty Botch | 600-800 ml | Apply as a full cover spray in 500-1000L of water per hectare at pink bud or before onset of the diseases at 7 to10-day intervals up to a maximum of 3 sprays per season. Use higher rate and apply at shorter interval under heavy disease pressure. |

**Contact: Robert Mazuru / Craig Chittenden / Dzingo Mafuvadze / Golden Zenda**

**Mobile: 0712 204 946 - 7 / 0773 212 051**

**Email: [mazuru@pivotalagro.com](mailto:mazuru@pivotalagro.com) / [craig@pivotalagro.com](mailto:craig@pivotalagro.com) / [dzingo@pivotalagro.com](mailto:dzingo@pivotalagro.com)**

