



# THIARA®250 WG

**A SYSTEMIC INSECTICIDE WITH STOMACH AND CONTACT ACTION FOR CONTROL OF VARIOUS SUCKING INSECT PESTS ON VARIOUS CROPS.**

THIARA®250 WG can be applied as a soil drench, foliar spray and through micro-jet or drip irrigation. Apply first at planting or before onset of pest infestation and repeat 14-28 days later. Maximum of two applications per season.

## **APPLICATION RATES & METHODS:**

| <b>CROP</b>                                 | <b>PEST</b>                                       | <b>RATE/100L<br/>OR<br/>Per Ha</b> | <b>Water Vol.<br/>(L/ha)</b> | <b>APPLICATION METHOD</b>                                                                                                                                                                   | <b>PHI<br/>(Days)</b> |
|---------------------------------------------|---------------------------------------------------|------------------------------------|------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| <b>Brassicas</b>                            | <i>Aphids</i>                                     | 200g/ha                            | 1000                         | Soil drench                                                                                                                                                                                 | 50                    |
| <b>Blueberries</b>                          | <i>Fruit Weevil</i>                               | 250g/ha                            | 500                          | Soil drench                                                                                                                                                                                 | 60                    |
| <b>Capsicums</b>                            | <i>Stinkbugs</i><br><i>Weevils</i>                | 250g/ha                            | 750                          | Soil drench                                                                                                                                                                                 | 30                    |
| <b>Citrus</b>                               | <i>Red scale</i><br><i>Psylla</i>                 | 250g/ha                            | 1000                         | Soil drench or Irrigation                                                                                                                                                                   | 30                    |
| <b>Cucurbits</b>                            | <i>Aphids</i>                                     | 200g/ha                            | 1000                         | Soil drench                                                                                                                                                                                 | 3                     |
| <b>Macadamia</b>                            | <i>Stinkbugs</i>                                  | 8g/tree                            | 500                          | Soil drench or Irrigation                                                                                                                                                                   | 90                    |
| <b>Ornamentals</b>                          | <i>Aphids</i><br><i>Stinkbugs</i>                 | 250g/ha                            | 750                          | Soil drench or Irrigation                                                                                                                                                                   | 14                    |
| <b>Pome Fruit</b><br><b>Stone Fruit</b>     | <i>Wooly Aphids</i>                               | 250g/ha                            | 500                          | Soil drench or Irrigation                                                                                                                                                                   | 60                    |
| <b>Potato</b>                               | <i>Aphids</i><br><i>Snout beetle</i>              | 250g/ha                            | 500                          | Soil drench or Foliar<br>Spray                                                                                                                                                              | 60                    |
| <b>Tomato</b>                               | <i>Whitely</i><br><i>Thrips</i>                   | 250g/ha                            | 500                          | Soil drench                                                                                                                                                                                 | 7                     |
| <b>Root<br/>Vegetables</b>                  | <i>Aphids</i><br><i>Hoppers</i>                   | 200g/ha                            | 1000                         | Soil drench                                                                                                                                                                                 | 30                    |
| <b>Tobacco Float<br/>Seedbed</b>            | <i>Aphids</i>                                     | 3 g                                | 100                          | Apply mixture into Float seedbed water at 7 weeks after sowing. Repeat application 21 days later if necessary.                                                                              | N/A                   |
| <b>Tobacco<br/>Conventional<br/>Seedbed</b> | <i>Aphids</i>                                     | 25 g                               | 100                          | Apply mixture as a soil drench at 1.5L/m <sup>2</sup> before sowing                                                                                                                         | N/A                   |
| <b>Tobacco Lands</b>                        | <i>Aphids</i><br><i>False</i><br><i>Wireworm</i>  | 125 g                              | 450                          | Apply 30 ml of mixture in the planting hole at time of planting                                                                                                                             | N/A                   |
| <b>Sugarcane</b>                            | <i>Yellow</i><br><i>Sugarcane</i><br><i>Aphid</i> | 870 g                              | 200 -300                     | <b>Plant cane</b><br>Apply as in-furrow band application at planting before covering.<br><b>Ratoon cane</b><br>Apply 7 – 30 days after harvesting as a band application over the cane rows. | 180                   |

**Contact: Robert Mazuru / Craig Chittenden / Dzingo Mafuvadze**

**Mobile: 0712 204 946 - 7 / 0773 212 051**

**Email: [mazuru@pivotalagro.com](mailto:mazuru@pivotalagro.com) / [craig@pivotalagro.com](mailto:craig@pivotalagro.com) / [dzingo@pivotalagro.com](mailto:dzingo@pivotalagro.com) / [golden@pivotalagro.com](mailto:golden@pivotalagro.com)**